

dispose to temporary paralysis, as it is met with so often in this state, but it may be questioned whether paralysis would sufficiently account for rupture of the bladder from over-distension. As an example of the persistence of spasm, a case may be mentioned here which occurred while I acted as house-surgeon in the Glasgow Royal Infirmary, of a man who was admitted suffering from retention of urine from spasmodic stricture, consequent on excessive indulgence in alcohol. I passed a No. 10 catheter easily, and withdrew ninety ounces of urine, carefully measured. The only result was paralysis of the bladder for some time. The semi-unconsciousness in the present case seems to have been continuous from the Saturday till death, the alcoholic stupor gradually merging into the drowsiness and coma of uræmic poisoning.

N.B. The preventability of rupture in this case is obviously beyond comment.

Glasgow.

A GRAVE COMPLICATION OF TYPHOID FEVER.

BY C. F. MAUNDER,
SURGEON TO THE LONDON HOSPITAL.

TYPHOID FEVER having just lately attracted a good deal of attention from the profession, two cases associated with hernia which have come under my observation will have some interest for it.

CASE 1 is that of a feeble old gentleman seen by me in consultation with Dr. Gillies, after three or four days' illness. He had been the subject of what proved to be a direct inguinal hernia of the right side, which the doctor thought he had partially reduced by taxis; but vomiting persisted, and was now stercoraceous. Aperient medicine had been rejected by the stomach, and there was constipation. His illness apparently began with sudden pain in the right inguinal region. I explored the inguinal swelling, and disclosed an old hernial sac, empty, with a very narrow neck indeed, and loaded with subperitoneal fat. Some days subsequently I heard from Dr. Gillies that our patient was undoubtedly the subject of typhoid fever, and, later still, that he had recovered, in spite of the surgeon, but much to the credit of the physician, as I think.

Of course, in the early stage of possible fever, with no special symptom to guide, obstinate and stercoraceous vomiting with a history of hernia and a swelling in a hernial region demand an exploratory operation. It is preferable to perform such an operation unnecessarily than to risk the possibility of death from strangulated hernia.

CASE 2.—A young man who had been ailing for many days was the subject of an inguinal hernia of the left side. The belly was tympanitic and tender, and he kept the left thigh flexed on the pelvis to relieve discomfort in the inguinal region. Handling the inguinal swelling caused pain. Constipation with nausea and sickness existed. His dull, listless aspect led me to look for a rose rash, and three or four spots were found. I now obtained the aid of Dr. Down, who pronounced the case to be one of typhoid fever. The question of operation was no longer entertained, and the progress of the case showed it would have been superfluous.

In this latter instance, but not in the former, the duration of the illness associated with rose rash materially aided the diagnosis; while, on the other hand, the flexed left thigh and tender bubonocoele tended to throw the observer off his guard.

Queen Anne-street.

ON THE CURE OF A LARGE VARICOSE ULCER BY TANNING WITH BABOOL BARK.

BY SURGEON J. ROCHE, M.D., BOMBAY ARMY,
CIVIL SURGEON, ROTREE, INDIA.

CAPT. F—— commanded one of the Indus Steam Flotilla Company's steamers which ply on the river Indus between Rotree, in Scinde, and Mooltan, in the Punjab. His habits were those of a free liver at table, without being intemperate. His weight, I should say, was nearly sixteen stone.

He was under my care for treatment in March, 1871, with a large varicose ulcer in the front and outer part of the leg. For many years he had been inconvenienced by varicose veins in the same leg, extending up the thigh to the groin, and, to relieve the consequent sense of weight and distension in the limb, he had worn an elastic stocking. The varicosity had affected the minutest veins, so as to change the colour of the integument in patches of black, brown, or red.

A few months before his application to me, when he was returning to his ship at night, he fell across the iron rails which are close to the river to permit the railway carriages to be run in proximity to the steamers whilst discharging cargo. He paid little attention to the contusion and abrasion which resulted, and proceeded to Mooltan as usual in command of his ship.

The great difficulty in navigating the river Indus when it is low in the cold weather, and to avoid the shifting sands in the bed of the river, necessitate the commander's unremitting attention from early dawn to dark every day. The standing posture which Capt. F—— was constrained to maintain quickly inflamed his leg, and he was forced to discard the use of the elastic stocking. The voyage to Mooltan in the cold weather is tedious in the extreme, and, though a distance of 500 miles, has been known to occupy six weeks, from the steamer having rested on sands. Towards the end of March, as the river rose, the navigation became easier, and Capt. F—— was able to give intervals of rest to his painful limb on the return voyage.

When he applied to me there was an inflamed and sloughing ulcer on the front and outer part of the leg. The central part of the ulcer looked like meat which had been exposed to the sun till decomposition commenced, and from it was issuing an ichorous fluid. The circumferential portion was covered by a shining and translucent epidermis, which was at once perceived to be dead by the vitreous and almost crystalline character it gave to the tissues underneath. The foot and leg were much greater in girth than their fellows on the other side.

The treatment prescribed embraced rest, elevation of the limb, poultices, a cholagogue, and abstemiousness. In a few days the sloughy surface was cleared, and there was left a large sore, red, angry, and painful—very like one consequent on a burn. Carron oil was used as dressing, and seemed to answer well for a few days. The angry appearance continued, if not increased. Bread-and-water poultices (over which a lotion, containing forty grains each of tannic acid and alum, with a drachm of tincture of belladonna, to ten ounces of water, was poured) were used. The ulcer soon lost its irritable character, but it became pale and indolent. The new epidermis which was forming was quite unhealthy. A dressing composed of half-an-ounce each of castor and turpentine oil to six ounces of linseed oil was tried, but the patient could not bear it longer than about two hours, when he said he felt as if a red-hot iron was fixed on his leg. Zinc lotion (a drachm and a half to eight ounces of water) was used. Tannic acid and alum lotion was, after some days, again had recourse to; the pledget of lint wrung out, which was covered over with a poultice or a piece of oiled silk at night, as a less or greater temperature was desired to be maintained. I am of opinion that the temperature maintained by a poultice is much less than that kept up by a piece of oiled silk. Strapping was tried when the granulations were large and weak. Bandaging the foot and leg to a short distance from the raw surface was done regularly. Under each form of dressing—the sedative when angry and irritable, the stimulating and supporting when indolent, cold lotions and water-dressing when inflamed, and poultices when sloughy—the ulcer did temporarily well, and many times changed its character. It frequently diminished in size to about two inches and a half in diameter, and as often enlarged to its original dimensions of nearly six inches. This intractable feature had almost completely exhausted the patience of the sufferer and myself, and lasted nearly two months. The internal medicines exhibited comprised sedatives, tonics, purgatives, and iodide of potassium.

Skin-grafting was often thought of, but I had not much faith in it, as some of the ulcer was more than once healed over by skin or its analogue, and which as often had died off. It occurred to me to get the ulcer a little weak with water-dressing, and literally tan it. I asked Capt. F—— to